

Comment on “Two Phase Transitions in the Fully frustrated XY Model”

The fully frustrated two-dimensional XY model (FFXYM) has been the source of many recent studies, and controversies [1–6]. The reason for this interest is that this model appears to have rather unconventional critical properties. Furthermore, this model has a clear experimental representation in Josephson junction arrays in magnetic fields. The model has $U(1)$ and $Z(2)$ symmetries, associated to its phase and chiral, or Ising, degrees of freedom, respectively. These symmetries are intertwined in purely thermodynamic quantities, like the helicity modulus, Υ . The possibility of defining a new universality class to describe the properties of the model appears to depend on having or not both symmetries breaking at the same temperature. Answering this question conclusively has turned out to be very hard since both transitions seem to occur either very close or may be on top of each other.

In a recent Letter [1] Olsson has argued that (i) both transitions take place at different temperatures, (ii) that the jump in Υ_{FFXYM} is of the standard unfrustrated 2-D XY Berezinskii-Kosterlitz-Thouless (BKT) type and (iii) that the chiral exponent $\nu_{Z(2)} = 1$, as in the standard 2-D Ising model, contradicting a large number of previous papers that obtained $\nu_{Z(2)} < 1$ [2,4–6]. Here we argue that the analysis given to reach these conclusions has flaws and the conclusions are not consistent with the evidence presented. Furthermore, we present new data of a more extensive MC calculation that gives $\nu_{Z(2)} < 1$.

(i-ii) The conclusion of having two different transition temperatures is based in part on the assumption that $\Upsilon_{FFXYM}(T_c) \equiv \Upsilon(T_{BKT})$. This translates into setting the exponent $\eta = \frac{1}{4}$ exactly. A very important element missing in the analysis of [1] is that there is no explicit mention of any type of error analysis carried out in the calculations. It is well known that this type of analysis is crucial in reliably determining exponents in XY-like models via MC calculations [7]. Then the claim that the two critical temperatures are different an equal to $T_{BKT} = 0.446$ and $T_{Z(2)} = 0.452$, which differ by about 3%, without quoting any error bars appears to be overly optimistic. A similar analysis, including errors, was done in [3], where they instead find a *non-universal* jump of 0.22(2). In contrast to [1,3] we have carried out an extensive explicit calculation of the gauge invariant zero momentum $U(1)$ correlation function and we found *instead* $\eta \approx \frac{1}{5}$ [4]. Of significant importance is that this result has been successfully compared with experimental results in JJA [6].

(iii) Olsson calculated the chiral correlation function and then its coherence length $\xi_{Z(2)}$, from which he got $\nu_{Z(2)} = 1$. One important drawback of this calculation is that the data analyzed was taken about 15% away from $T_{Z(2)} = 0.452$. In [4] we concentrated in the calculation of the $U(1)$ correlation functions and less so in the chiral one. We have now extended our analysis of

the zero momentum chiral correlation function, and the $\xi_{Z(2)}(T)$ data is shown in the table. We have carefully

TABLE I. Results for $\xi_{Z(2)}$ obtained from $g_{z(2)}(r)$ at different temperatures and lattice sizes.

T	L	$\xi_{Z(2)}$	NMCS	L	$\xi_{Z(2)}$	NMCS
			K			K
0.70	24	0.8217 (671)	160	32	0.9312 (92)	160
0.60	24	1.3341 (68)	160	32	1.5520 (530)	160
0.575	24	1.4318 (73)	160	32	1.7579 (62)	160
0.55	24	1.5386 (43)	160	32	2.1278 (26)	160
0.525	24	2.3948 (37)	175	32	2.9035 (101)	175
0.50	48	3.8496 (37)	175	64	4.4840 (38)	150
0.49	48	4.3700 (36)	225	64	5.9480 (37)	225
0.48	48	5.3916 (20)	225	64	7.2132 (432)	225
0.47	48	6.4206 (82)	225	64	9.8993 (586)	225
0.46	—	—	—	128	20.2928 (464)	175

analyzed the data, together with an error analysis, as done in [4,7], in an extended temperature range, and we obtained $\nu_{Z(2)} = 0.898(3)$ and $T_{Z(2)} = 0.4511(10)$. The critical exponent is in good agreement with previous calculations while the critical temperature is close to the one obtained in [1]. Furthermore, recent experiments in superconducting wire networks [8], where the order parameter fluctuations are important, have also found that the chiral exponent $\nu_{Z(2)} \neq 1$.

This work has been partially supported by grant NSF-DMR-95-21845 (JJV), and by DGAPA-UNAM grants IN-103294 and IN-100595 (GRS). The calculations were mostly done at DGSCA-UNAM.

Jorge V. José¹ and G. Ramírez-Santiago²

¹ Department of Physics and Center for Interdisciplinary Research on Complex Systems, Northeastern University, Boston MA, 02115.

² Instituto de Física, UNAM, P.O. Box 20-364, 01000 México D.F.

- [1] P. Olsson, Phys. Rev. Lett. **75**, 2758 (1995).
- [2] e.g. J. Lee, et al. Phys. Rev. B **43**, 11531 (1991); E. Granato and M.P. Nightingale, *ibid* **48**, 7438 (1993); J.-R. Lee, *ibid* **49**, 3317 (1994).
- [3] S. Lee and K. C. Lee Phys. Rev. B **49**, 15184 (1994).
- [4] G. Ramírez-Santiago and J.V. José, Phys. Rev. Lett. **68**, 1224 (1992); Phys. Rev. B **49**, 9567 (1994).
- [5] M.P. Nightingale, et al. Phys. Rev. B **52**, 7402 (1995).
- [6] J. V. José, et al. Physica B+C **194-196**, 1671 (1994).
- [7] R. Gupta and C. F. Baillie, Phys. Rev. **B45**, 2883 (1992).
- [8] X. S. Ling et al. Phys. Rev. Lett. **76**, 2989 (1996).